


"ALEXANDRU IOAN CUZA" UNIVERSITY OF IASI
FACULTY OF GEOGRAPHY AND GEOLOGY
FIELD OF ENVIRONMENTAL SCIENCE
DOCTORAL SCHOOL OF GEOSCIENCES



**THE INFLUENCE OF SALINE AEROSOLS ON HUMAN HEALTH AND
PERFORMANCE**

PHD THESIS SUMMARY

Scientific coordinator:

Prof. univ. dr. ION SANDU

PhD student:

MIHAELA-ORLANDA ANTONOVICI (CAS. MUNTEANU)

IASI, 2023

Abstract

The paper presents the experimental data obtained on two groups of students, one witness, in two normal classrooms and respectively in the outdoor gymnasium and another in the same type of rooms, but with halochamber regime, differentiated on gender and age, including children born in 2006 and 2007, at the age with the highest growth rate. Starting from the fact that the solions reformulated *in situ* act on the body through inhalation and absorption at the skin level, it offers a remedy for many conditions, as halotherapy is a complementary and / or alternative method of protecting the airways and improving the morpho-structural characteristics of the skin of the human body. An original system was used to process the experimental data regarding the evolution of the characteristics studied, expressed as a percentage increase, allowed a good highlighting of the effect of saline aerosols/solions on the morpho-structural type biometric characteristics (weight, height, thoracic perimeter), pulse and blood pressure in children, depending on age and gender. It has been shown that girls interact more strongly with solions, and register higher values for most of the studied parameters than boys.

Saline aerosols in the form of nano dispersions in gaseous environments behave, as far as concentration levels and lifetime, in the form of trimodal Gaussian distributions: *fine Aitken particles* (below 50 μm), *medium* (between 50 and 500 μm) and *coarse or large* (between 500 and 1000 μm) respectively. The fine and medium ones have the stability over time depending on the active surface of the porous granules of the source of generation (diaphragm), the shape (cylindrical or parallelepiped diaphragm), size and structural profile of the diaphragm, number of generation centers, particle size and position of superficial fluorescent microcrystallites. superficial (external surface and that of the pores of the granules), the porosity of the granules, the levels of the climatic parameters and other characteristics of the gaseous environment of the halochamber.

The works elaborated in the last years within our group have experimentally demonstrated that the solion-type hydrated nano-aerosols, as such or doped with other halo-salts, in addition to the ability to prevent and treat diseases cardio-respiratory, osteo-muscular and psychomotor, have at certain levels of concentration beneficial effects on the development of children, in increasing the athletic performance in young sportsmen and in improving the resistance of the persons who carry out activities with intense physical effort, in severe or high performance work regimes.

For people working in severe conditions (eg. welders), the aerial metal smoke resulting from the welding process can be inhaled by welders, being used in studies, respiratory filters (FFR), which stop the particles from entering the airways of the workers.

Research has been carried out on the effect they have on relative humidity, particles composed of sodium chloride and sodium sulfate, thus it was investigated how the particle size influences the efflorescence rate and which of them nucleate faster, when the NaCl particles are larger than 70 nm, the ERH decreases with decreasing dry particle sizes.

Therefore, in the particle size distribution of saline aerosols (Aitken nanoparticles of NaCl), two subgroups are differentiated: *anhydrous* and *hydrated* ones, called *solions*. The first are non-symmetrical, poorly aquatized nanopolyhedrons, which by the anhydrizing effect of the membrane systems will chronicize the affections, while the *solions* (continuously/ dynamically reformed at atmospheric humidities greater than 85%), with a movable *glomerular* structure at the level of a tissue *in vivo* through deliquescence will facilitate the directing of the ions to the specific channels, stimulating the membrane biochemical processes.

In other words, solions have good bioactive compatibility with living tissues, activating a number of membrane processes *in vivo*. Moreover, they, in addition to creating clean air

environments, allow, as previously mentioned, the prevention/treatment of certain diseases and to improve the physical performance of human subjects.

Aerosol therapy (inhalation therapy) is a field of science that deals with the methods of treatment and prevention of airway disease, the efficiency of the treatment depends on the size distribution of the aerosol particles, their physical properties (density, shape, hygroscopicity, morphology).

In connection with these nanoparticles, a series of patented artificial halochambers were developed, which were experienced in the teaching activities with the students. With their help it has been shown that solions active at a certain level of concentration have a beneficial influence on the development of the young, with the female having a higher receptivity regarding the effect of improving human performances.

The paper presents the influence of solions on the development of children, aged between 12 and 15 years, in terms of the evolution of body weight, height, chest perimeter or chest thickness, for a period of 15 months. In this respect, the influence of the active residence time in the halochamber is studied, which is ten times higher than in other studies of the group.

The characteristics of the solions during the rest periods, 50 minutes after the start and the end of the classes, were determined using two particle counters type SIBATA GT 321 and Dust Trok, an Air Ion Counter type ionometer and by the method of differential conductometry, using a patented device within our group. Also, the evolution in time and work stages of the following parameters registering the main climatic factors in the classrooms were monitored, namely: temperature ($^{\circ}\text{C}$), air humidity (% UR), atmospheric pressure (mmHg) and lighting (lx), using specific laboratory devices.

The study used a series of generators with diaphragms containing porous granules of NaCl, obtained by recrystallization, at 50... 60 $^{\circ}\text{C}$, from supersaturated solutions. Four generators were placed at the windows of two classrooms with a volume of 126 cubic meters (7 \times 6 \times 3) and six in the 320m³ (10 \times 8 \times 4) gyms. In order to achieve an optimal level in solions, the generators were started about 60 minutes before the start of class hours. After 50 minutes of operation, the average level of solions was determined, and then determined again at the end of the classes. For the Physical Education classes, there was a progressive program of exercises, training and games over the period of 50 minutes, twice a week, distributed on even time intervals. For the two groups of activities (normal lessons in the classroom and physical education in the school gym) there were two groups of students who performed in parallel, one group in salt rooms and another in normal classrooms or gym, using the same work schedule and the respective time intervals. The students were monitored throughout the school year, twice a week regarding the evolution of morphostructural and biometric characteristics. For each work program, the level reached by saline aerosols/solions was correlated with the average data of the biometric characteristics of the students, separated on gender.

The experiments were performed on two groups of students (girls and boys) from classes VI and VII, for each age category, noted by the year of birth (2006 and 2007): the reference group for outdoor or gym applications, made of the group of girls noted F0i200X (X = 6 and 7), with student i, having the year of birth 200X and the group of boys noted B0i200X, with student i, having the year of birth 200X and respectively the halochamber group formed by the group of girls noted Fhi200X having student i, with the year of birth 200X and the group of boys noted Bhi200X with student i, having the year of birth 200X, where $i = 1, 2, 3, \dots$, represents the number of the student in the group, each group with 7 to 25 students.

In order to select the students for the two experimental groups (from the halochamber and from the open air), a series of data was used according to a standard form that includes information regarding the family living conditions, the family medical history (hemophilia, diabetes mellitus, HTA, parasites, heart disease, cancer, rheumatic diseases, TB), personal physiological, pathological history (eruptive diseases, whooping cough, hepatitis, gastritis, cardiovascular, immunological, metabolic and endocrine disorders) .

Within the morphometric and functional biometric examination, in order to determine the level of physical development of the subjects, a series of measurements were made, namely:

- the weight, which was made with the help of the electronic scale and was expressed in kilograms;
- stature (height), which was determined with the staturometer, being expressed in centimeters;
- the thoracic perimeter, respectively the thickness of the thorax, was measured with a metric band (100cm), recording two dimensions, with the thoracic perimeter in respiratory rest (the moment between inspiration and expiration) and respectively the thoracic perimeter in maximum inspiration.

In order to facilitate the interpretation of the experimental results, the percentage increase of the characteristic over time was used graphically. For this purpose, the biometric data were determined periodically, at 3, 6, 9, 12, 15, 18, 21 and 24 months respectively for establishing the evolution of the subjects' weight and height, and for highlighting the increase of the thoracic perimeter, at 3, 6, 9, 12, 15, 18 and respectively 21 months. Based on the recorded data, a primary table was prepared. Then the average of the values for the same period evaluated for each group of students, respectively the group of girls, noted F0m200X and the group of boys, noted B0m200X, where m represents the arithmetic mean, and X - values 6 or 7, corresponding to the year of birth. . After this, the percentage (%) was calculated with the evolution of the characteristic C, compared to the start time of the experiment (C0), in relation to the subsequent readings (Cj), the period of determination being denoted by j, which represents the month of the year of readings, using relationship:

$$C (\%) F0m200X \text{ or } C (\%) B0m200X = 100 \times (Cj - C0) / C0, (1)$$

respectively:

$$C (\%) Fhm200X \text{ or } C (\%) Bhm200X = 100 \times (Cj - C0) / C0; (2)$$

Finally, the graphs $C (\%) = f(t)$ are drawn, the time being in months, for the average values of the girls and boys, in the two different working conditions, taken separately (outdoors, noted in the lower right indicates "0" and halochamber, noted down right indicates "h"). Each biometric characteristic, taken in the study, was represented in the same coordinate system, having in order C (%) and in the abscissa time in months. Based on these graphs, the evolution of each group of students was appreciated and the influence of saline aerosols and reformulated solions in situ on the characteristics studied was established.

Since the dynamics of aerosol emissions from halochambers is influenced by the environmental conditions, humidity, temperature, pressure and lighting, the determinations were made keeping the values of the parameters of the pumped air constant and monitoring the microclimate formed in the salt room. The main factor for improving and maintaining student health in our case is the sodium chloride solution, made up of nanoparticles with diameters ranging from 2 to 5 μm , structurally reformed from Aitken particles in a humid environment ($\geq 65\% \text{ RH}$).

The experimental data obtained on the groups of students, one witness, who worked in the outdoor sports hours or on the sports field (in good weather)/the gymnasium (in bad weather), respectively for the other teaching hours in two classes without aerosols and two halochamber type classes (classrooms equipped with dynamic aerosol/solion generators), the students being

differentiated by the two genders and ages, including children born in 2006 and 2007, and the time spent sitting/working in the halochamber being ten times more than the one studied before the following conclusions can be made:

- saline aerosols and solions reformulated in situ in humid environment respectively act on the body through inhalation and absorption at the skin level, which according to the literature offers a remedy for many conditions, halotherapy being a complementary and/or alternative method of airway protection, but and improving morpho-structural/biometric characteristics of the human body;
- the system of processing the experimental data regarding the evolution of the three morpho-structural/biometric characteristics (body weight, height and circumference of the thoracic perimeter), expressed as a percentage increase, allowed a good highlighting of the effect of saline aerosols/solions on the three characteristics taken in you study, by age and gender;
- it can be observed that for the girls born in 2006 the two curves of weight gain are strongly separated, the curve corresponding to the halochamber is higher in the first 18 months than the one in the witness room. In contrast, for boys the increase in weight for the two curves (from the halochamber and from the witness room) have parallel and similar values and close evolution rates, but after 18 months they are above the development curves in girls; this effect is due to the fact that boys have a higher development rate than girls, while girls are more vulnerable to the influence of solions;
- the same effect is found for students born in 2007, with the difference that from the beginning the evolution of weight for boys is above that corresponding to girls;
- in terms of height, both for students born in 2006 and 2007, the percentage growth rate for girls is higher than for boys;
- the third characteristic, the circumference of the thoracic perimeter, presents for the students born in 2006 a higher percentage growth rate than for the girls, and for those born in 2007 it is located between curve related to the halochamber and that of the witness room corresponding to girls;
- the increase of the average dimensions of the thoracic perimeter dimensions recorded in respiratory rest and respectively in maximum inspiration (*inspire*) under the influence of saline aerosols/solions, for the largest age series, presents a lower growth rate in girls than in boys, a phenomenon considered normal;
- on the other hand, for the younger series, the same evolution of the size of the thoracic perimeter registers a higher growth rate in girls than in boys, and in the first 12 months the values recorded in inspiration, both in the case of the witness and in the halochamber groups are very close, so that later, the average size of the thoracic perimeter will increase more strongly in girls, than in boys, an evolution that can be explained by the genetic diversity and the existence of large differences in the quality of life of children in the study area (rural, mountain with oil and forestry industry);
- Class and physical education classes in schools in an environment with saline aerosols / solions can help the proper functioning of the body, they have a positive impact on school activities and sports competitions of students, implicitly on human performance;
- the experimental data processing system regarding the evolution of the characteristics studied, expressed by percentage increase, allowed a good highlighting of the effect of saline aerosols / solions on the biometric characteristics of morpho-structural type (weight,

height, thoracic perimeter) in children, these have been studied and exposed in a previous article according to age and sex;

- the existing microclimate in saline has a major contribution on the human body, numerous studies conducted so far, have registered favorable results only in the prevention and treatment of non-chronic respiratory diseases, without paying attention to the impact on intellectual and physical performance of students, grouped by age, sex, control and experimental group. The research of our team targeted high school students, on whom saline aerosols had a special influence;
- saline aerosols and solions reformulated in situ act on the body by inhalation and absorption in the skin, which according to the literature offers a remedy for many diseases, halotherapy being a complementary and / or alternative method of airway protection and improvement of the morpho-structural characteristics of the human organism;
- the correct and systemic practice of movement games in an environment with saline aeroanions / solions contributes substantially to the development of students' resistance to diseases, to a better functioning of the body and to obtaining sports performances;
- the students who study in classes held in schools in an environment with saline aerosols / solions can help the proper functioning of the body, they have a positive impact on school activities and sports competitions of students;

The experimental data processing system regarding the evolution of the characteristics studied, allowed a good highlighting of the effect of saline aerosols / solions on some functional biometric characteristics in children, depending on age and sex.